

A man was diagnosed with acute radiation sickness. During laboratory testing, a sharp decrease in serotonin in platelets was detected. The metabolism of which substance caused these indicators?

Tyrosine

Phenylalanine

5-hydroxytryptophan

Serine

Histidine

One of the pathogenetic links in the development of radiation sickness is the intensification of free radical oxidation processes. What substances are the main source of free radical formation?

Proteins

Lipids

Carbohydrates

Water

Hormones

The patient was diagnosed with the bone marrow form of acute radiation sickness. Which of the following hematological symptoms will be observed during the flare-up period?

Left shift of the leukocyte formula

Relative lymphocytosis

Relative lymphopenia

Pancytopenia

Erythrocytosis

As a result of radiation, hematopoietic stem cells were damaged.
The formation of which connective tissue cells will be disrupted?

Melanocytes

Pericytes

Adipocytes

Macrophages

Fibroblasts

After radiation exposure, the doctor recommended that the patient increase the content of vegetable oils in the diet - sources of polyene fatty acids. Name the acid that contains three double bonds:

Arachidonic

Myristic

Oleic

Lauric

Linolenic

As a result of radiation, hematopoietic stem cells were damaged.
The formation of which connective tissue cells will be disrupted?

Melanocytes

Pericytes

Macrophages

Fibroblasts

Adipocytes